School of Architecture, Land and Environmental Sciences

Articles

2006

The impact of park trees on microclimate in urban areas

Georgi, Neratzia Julia

Springer Verlag

http://hdl.handle.net/11728/7503

Downloaded from HEPHAESTUS Repository, Neapolis University institutional repository

The impact of park trees on microclimate in urban areas

N. J. Georgi · K. Zafiriadis Published online: 30 May 2006

c_Springer Science+Business Media, LLC 2006

Abstract Trees in cities have an important positive effect on people's lives. One such positive effect is the amelioration of microclimate. The aim of this research is to assess the correlation between parameters that affect the microclimate of parks during the summer. We measured air temperature, relative humidity and solar radiation in the sun and shade of the trees in urban parks in Thessaloniki, Greece. The results indicate that: the air (A) temperature reduction percentage (dTair%), air relative humidity increase (dTRh%), discontent index reduction percentage (DI%) (cooling effect) and solar radiation (L) percentage that passes through the trees' foliage to their shade creates an exponential function of dA% = a.e-bL. These functions are also applicable to the limiting variation values of the parameters. If we use L = 0 (meaning Lightsh = 0, which is the case for an extremely dense tree), then the values that we expect from this particular parameter are the maximum possible. These maximum values are a characteristic feature of the parameter variation for this particular research area. These maximum values for the trees in the parks of Thessaloniki are: maxdTair% $\approx 24\%$, maxdRh% $\approx 41\%$ and maxdDI% $\approx 16\%$.

 $\textbf{Keywords} \ \text{Air temperature} \cdot \text{Relative humidity} \cdot \text{Solar radiation} \cdot \text{Thermal comfort} \cdot \text{Shade trees}$